

Tender Heart High School, Sec.-33B chd.

class: VIII

Teacher: Deepa

Subject: Mathematics

Topic: Chapter - : Sets

In this week, we will cover the following topics

- Subsets
- Complement of Set
- Superset
- proper subset
- Universal Set
- Union / Intersection

Subsets: If A and B are two sets given in such a way that every element of A is in B, then we say that A is a subset of B, $\Rightarrow A \subseteq B$
 \Rightarrow If $A \subseteq B$, then every element ($x \in A$) also in $x \in B$

If $A \subseteq B$ then B is called superset.

- There are 2^n subsets of a given set.
- \emptyset is the subset of every element (Set.)
- Every set is a subset of itself.

Proper Subset: Let A and B are two set, then A is called proper subset of B if all the element of A is present in B, but there is at least one element in B which is not in A.

$$\text{i.e } \{1, 2\} \subset \{1, 2, 3, 4\}$$

so set $\{1, 2\}$ is a proper subset of $\{1, 2, 3, 4\}$

Note: \emptyset has no proper subset.

for e.g

Ques 1. Write all the subsets of $B = \{a, b, c\}$

Soln: $\emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{b, c\}, \{a, b, c\}$

There are $2^3 = 8$, Subsets of the given set.

Class: VIII

Subject: Mathematics

Topic: Chapter-1: Sets

classmate

Date _____

Page _____

2

Teacher: Deptt.

Quest. 2 Write the proper subset of set $A = \{1, 2, 3\}$
Solt: $\emptyset, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}$
There are $(2^n - 1) = (8 - 1) = 7$ proper subset
of the given set A

Operations on Sets:

1. Union of Sets: It is written as $A \cup B$, means
containing all the elements of
A as well as of B.
for eg:

If $A = \{1, 2, 3, 4\}$ and $B = \{4, 5, 6, 7\}$
then $A \cup B = \{1, 2, 3, 4, 5, 6, 7\}$

2. Intersection of sets: It is written as $A \cap B$, means
set of elements common in
both the given sets.
for eg:

If $A = \{1, 2, 3, 4\}$ and $B = \{4, 5, 6, 7\}$
then $A \cap B = \{4\}$

only the common elements.

* If the sets are disjoint, then $A \cap B = \emptyset$

3. Difference of sets: for any two sets A and B,
the difference $A - B$ is the
set of all the elements of A which are
not in B.

for eg:

If $A = \{1, 2, 3, 4\}$ and $B = \{4, 5, 6, 7\}$
then $A - B = \{1, 2, 3\}$

Class: VIII

Subject: Mathematics

Topic: Chapter- 1: Sets

classmate

Date _____

Page 3

Teacher: Deepika

4. Complement of a set: Let U or \mathcal{E} be an universal set and $A \subseteq \mathcal{E}$ then complement of A i.e A^c or A' is the set of elements of U not in A .

Let the universal set be $U = \{1, 2, 3, 4, 5, 6, 7, 8\}$ and $A = \{4, 5, 6\}$ then $A^c = \{1, 2, 3, 7, 8\}$

Points to remember:

(i) $\emptyset' = U$ and $U' = \emptyset$

* Complement of an empty set is Universal set.

* Complement of a Universal set is an empty set.

(ii) $A \cup A' = U$ and $A \cap A' = \emptyset$

* Union of a set and its complement is "Universal set"

* Intersection of a set and its complement is an "empty set"

